



TP53 gene

tumor protein p53

Normal Function

The *TP53* gene provides instructions for making a protein called tumor protein p53 (or p53). This protein acts as a tumor suppressor, which means that it regulates cell division by keeping cells from growing and dividing too fast or in an uncontrolled way.

The p53 protein is located in the nucleus of cells throughout the body, where it attaches (binds) directly to DNA. When the DNA in a cell becomes damaged by agents such as toxic chemicals, radiation, or ultraviolet (UV) rays from sunlight, this protein plays a critical role in determining whether the DNA will be repaired or the damaged cell will self-destruct (undergo apoptosis). If the DNA can be repaired, p53 activates other genes to fix the damage. If the DNA cannot be repaired, this protein prevents the cell from dividing and signals it to undergo apoptosis. By stopping cells with mutated or damaged DNA from dividing, p53 helps prevent the development of tumors.

Because p53 is essential for regulating cell division and preventing tumor formation, it has been nicknamed the "guardian of the genome."

Health Conditions Related to Genetic Changes

bladder cancer

Somatic *TP53* gene mutations have been found in some cases of bladder cancer. Most of these mutations change single amino acids in p53. The altered protein cannot bind to DNA, preventing it from effectively regulating cell growth and division. As a result, DNA damage accumulates in cells, which can allow them to grow and divide in an uncontrolled way to form a cancerous tumor. Mutations in the *TP53* gene may help predict whether bladder cancer will progress and spread to nearby tissues, and whether the disease will recur after treatment.

breast cancer

Inherited changes in the *TP53* gene greatly increase the risk of developing breast cancer, as well as several other forms of cancer, as part of a rare cancer syndrome called Li-Fraumeni syndrome (described below). These mutations are thought to account for only a small fraction of all breast cancer cases.

Noninherited (somatic) mutations in the *TP53* gene are much more common than inherited mutations, occurring in 20 to 40 percent of all breast cancers. These somatic mutations are acquired during a person's lifetime and are present only in

cells that become cancerous. The cancers associated with somatic mutations do not occur as part of a cancer syndrome. Most of these mutations change single protein building blocks (amino acids) in the p53 protein, which reduces or eliminates the protein's tumor suppressor function. Because the altered protein is less able to regulate cell growth and division, DNA damage can accumulate. This damage may contribute to the development of a cancerous tumor by allowing cells to grow and divide in an uncontrolled way.

Compared with breast cancers without *TP53* gene mutations, tumors with these genetic changes tend to have a poorer prognosis. They are more likely to be aggressive, to be resistant to treatment with certain anti-cancer drugs and radiation, and to come back (recur) after treatment.

cholangiocarcinoma

head and neck squamous cell carcinoma

Somatic mutations in the *TP53* gene have been found in nearly half of all head and neck squamous cell carcinomas (HNSCC). This type of cancerous tumor occurs in the moist lining of the mouth, nose, and throat. Most of the *TP53* gene mutations involved in HNSCC change single amino acids in p53; these changes impair the protein's function. Without functioning p53, DNA damage builds up in cells, and they can continue to divide without control, leading to tumor formation.

Li-Fraumeni syndrome

Although somatic mutations in the *TP53* gene are found in many types of cancer, Li-Fraumeni syndrome appears to be the only cancer syndrome associated with inherited mutations in this gene. This condition greatly increases the risk of developing several types of cancer, particularly in children and young adults. At least 140 different mutations in the *TP53* gene have been identified in individuals with Li-Fraumeni syndrome.

Many of the mutations associated with Li-Fraumeni syndrome change single amino acids in the part of the p53 protein that binds to DNA. Other mutations delete small amounts of DNA from the gene. Mutations in the *TP53* gene lead to a version of p53 that cannot regulate cell growth and division effectively. Specifically, the altered protein is unable to trigger apoptosis in cells with mutated or damaged DNA. As a result, DNA damage can accumulate in cells. Such cells may continue to divide in an uncontrolled way, leading to the growth of tumors.

ovarian cancer

Somatic *TP53* gene mutations are common in ovarian cancer, occurring in almost half of ovarian tumors. These mutations result in a p53 protein that is less able to control cell growth and division, contributing to the development of a cancerous tumor.

other cancers


Somatic mutations in the *TP53* gene are the most common genetic changes found in human cancer, occurring in about half of all cancers. In addition to the cancers described above, somatic *TP53* gene mutations have been identified in several types of brain tumor, colorectal cancer, liver cancer, lung cancer, a type of bone cancer called osteosarcoma, a cancer of muscle tissue called rhabdomyocarcinoma, and a cancer called adrenocortical carcinoma that affects the outer layer of the adrenal glands (small hormone-producing glands on top of each kidney).

Most *TP53* mutations change single amino acids in the p53 protein, which leads to the production of an altered version of the protein that cannot control cell growth and division effectively. As a result, cells can grow and divide in an unregulated way, which can lead to cancerous tumors.

Chromosomal Location

Cytogenetic Location: 17p13.1, which is the short (p) arm of chromosome 17 at position 13.1

Molecular Location: base pairs 7,668,402 to 7,687,550 on chromosome 17 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- antigen NY-CO-13
- cellular tumor antigen p53
- P53
- P53 tumor suppressor
- P53_HUMAN
- phosphoprotein p53
- transformation-related protein 53
- TRP53

- tumor protein p53 (Li-Fraumeni syndrome)
- tumor suppressor p53

Additional Information & Resources

Educational Resources

- Molecular Biology of the Cell (fourth edition, 2002): Cell-Cycle Progression is Blocked by DNA Damage and p53: DNA Damage Checkpoints
<https://www.ncbi.nlm.nih.gov/books/NBK26856/#A3240>
- Molecular Cell Biology (fourth edition, 2000): Mutations in p53 Abolish G1 Checkpoint Control
<https://www.ncbi.nlm.nih.gov/books/NBK21551/#A7158>
- National Cancer Institute: Genetics of Breast and Gynecologic Cancers (PDQ)
<https://www.cancer.gov/types/breast/hp/breast-ovarian-genetics-pdq>

GeneReviews

- Li-Fraumeni Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1311>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28TP53%5BTI%5D%29+OR+%28p53%5BTI%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+180+days%22%5Bdp%5D>

OMIM

- TUMOR PROTEIN p53
<http://omim.org/entry/191170>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/P53ID88.html>
- Cancer Genetics Web: TP53
<http://www.cancerindex.org/geneweb/TP53.htm>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=TP53%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=11998

- IARC TP53 Mutation Database
<http://p53.iarc.fr/>
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/7157>
- p53 Web Site (Thierry Soussi)
<http://p53.fr/>
- UniProt
<http://www.uniprot.org/uniprot/P04637>

Sources for This Summary

- Damineni S, Rao VR, Kumar S, Ravuri RR, Kagitha S, Dunna NR, Digumarthi R, Satti V. Germline mutations of TP53 gene in breast cancer. *Tumour Biol.* 2014 Sep;35(9):9219-27. doi: 10.1007/s13277-014-2176-6. Epub 2014 Jun 15.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24929325>
- GeneReview: Li-Fraumeni Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1311>
- Loyo M, Li RJ, Bettgowda C, Pickering CR, Frederick MJ, Myers JN, Agrawal N. Lessons learned from next-generation sequencing in head and neck cancer. *Head Neck.* 2013 Mar;35(3):454-63. doi: 10.1002/hed.23100. Epub 2012 Aug 21. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22907887>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3715072/>
- Masciari S, Dillon DA, Rath M, Robson M, Weitzel JN, Balmana J, Gruber SB, Ford JM, Euhus D, Lebensohn A, Telli M, Pochebit SM, Lypas G, Garber JE. Breast cancer phenotype in women with TP53 germline mutations: a Li-Fraumeni syndrome consortium effort. *Breast Cancer Res Treat.* 2012 Jun;133(3):1125-30. doi: 10.1007/s10549-012-1993-9. Epub 2012 Mar 4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22392042>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3709568/>
- Masica DL, Li S, Douville C, Manola J, Ferris RL, Burtneess B, Forastiere AA, Koch WM, Chung CH, Karchin R. Predicting survival in head and neck squamous cell carcinoma from TP53 mutation. *Hum Genet.* 2015 May;134(5):497-507. doi: 10.1007/s00439-014-1470-0. Epub 2014 Aug 10.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25108461>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4324386/>
- Merino D, Malkin D. p53 and hereditary cancer. *Subcell Biochem.* 2014;85:1-16. doi: 10.1007/978-94-017-9211-0_1. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25201186>
- National Cancer Institute: Genetics of Breast and Gynecologic Cancers (PDQ)
<https://www.cancer.gov/types/breast/hp/breast-ovarian-genetics-pdq>
- Olivier M, Hollstein M, Hainaut P. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol.* 2010 Jan;2(1):a001008. doi: 10.1101/cshperspect.a001008. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20182602>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2827900/>

- Olivier M, Langerød A, Carrieri P, Bergh J, Klaar S, Eyfjord J, Theillet C, Rodriguez C, Lidereau R, Bièche I, Varley J, Bignon Y, Uhrhammer N, Winqvist R, Jukkola-Vuorinen A, Niederacher D, Kato S, Ishioka C, Hainaut P, Børresen-Dale AL. The clinical value of somatic TP53 gene mutations in 1,794 patients with breast cancer. Clin Cancer Res. 2006 Feb 15;12(4):1157-67.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16489069>
- Ruijs MW, Verhoef S, Rookus MA, Pruntel R, van der Hout AH, Hogervorst FB, Kluijdt I, Sijmons RH, Aalfs CM, Wagner A, Ausems MG, Hoogerbrugge N, van Asperen CJ, Gomez Garcia EB, Meijers-Heijboer H, Ten Kate LP, Menko FH, van 't Veer LJ. TP53 germline mutation testing in 180 families suspected of Li-Fraumeni syndrome: mutation detection rate and relative frequency of cancers in different familial phenotypes. J Med Genet. 2010 Jun;47(6):421-8. doi: 10.1136/jmg.2009.073429.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20522432>
- Silwal-Pandit L, Vollan HK, Chin SF, Rueda OM, McKinney S, Osako T, Quigley DA, Kristensen VN, Aparicio S, Børresen-Dale AL, Caldas C, Langerød A. TP53 mutation spectrum in breast cancer is subtype specific and has distinct prognostic relevance. Clin Cancer Res. 2014 Jul 1;20(13):3569-80. doi: 10.1158/1078-0432.CCR-13-2943. Epub 2014 May 6. Erratum in: Clin Cancer Res. 2015 Mar 15;21(6):1502.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24803582>
- Toss A, Tomasello C, Razzaboni E, Contu G, Grandi G, Cagnacci A, Schilder RJ, Cortesi L. Hereditary ovarian cancer: not only BRCA 1 and 2 genes. Biomed Res Int. 2015;2015:341723. doi: 10.1155/2015/341723. Epub 2015 May 17. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/26075229>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4449870/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/TP53>

Reviewed: October 2015
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services